Author Index

Adaya Godoy, A., see Ladron de Guevara, O. 350
Ahel, M., see Czuczwa, J. 233
Alcazar, B., see Rubio, N. 312
Al-Thamir, W. K.
Gas chromatographic determination of water in crude oils 288
Amicosante, G.
——, Marinucci, M. C., Franceschini, N., Tizzani, M. I., Oliva, B. and Oratore, A.
Fractionation and characterization of two β-lactamases in *Citrobacter diversus* ULA-27 strain by chromatofocusing 366
Andersson, L., see Belew, M. 197
Andrikopoulos, N. K., see Moschidis, M. C. 363
Belew, M.
——, Yip, T.-T., Andersson, L. and Porath, J.
Interaction of proteins with immobilized Cu^{2+}. Quantitation of adsorption capacity, adsorption isotherms and equilibrium constants by frontal analysis 197
Bellama, J. M., see Eng, G. 1
Bergenhem, N.
——, Carlsson, U., Ellow, L.-E. and Rosén, S.
Rapid ion-exchange chromatography for preparative separation of proteins. III. Application to purification of bovine factor X and prothrombin 383
Bhide, S. V., see Talib, Y. Y. 373
Bours, J.
——, Dempewolf, J., Schaake, S. and Rink, H.
Determination of the electrophoretic order of 26 nucleotides by isotachophoresis 336
Bouthillier, F.
——, Carmel, M., Elhilali, M., Radwan, F. and Lehoux, J.-G.
Application of high-performance liquid chromatofocusing to the study of prostatic androgen receptors. Effect of stabilizing agents on the heterogeneity of receptor structure 171
Brieva, A., see Rubio, N. 312
Brinckman, F. E., see Eng, G. 1
Bystrický, S., see Strašák, M. 331
Cachet, Th.
——, Roets, E., Hoogmartens, J. and Vanderhaeghe, H.
Separation of novel derivatives from commercial erythromycin samples by thin-layer chromatography 343
Capri, S., see Marcomini, A. 243
Carlsson, U., see Bergenhem, N. 383
Carmel, M., see Bouthillier, F. 171
Charbonneau, C. F., see Lawrence, J. F. 379
Chen, A. J. C., see Hjertén, S. 47
Cortinas de Nava, C., see Ladron de Guevara, O. 350
Czuczwa, J.
——, Leuenberger, C., Tremp, J., Giger, W. and Ahel, M.
Determination of trace levels of phenol and cresols in rain by continuous liquid–liquid extraction and high-performance liquid chromatography 233
Dautrevaux, M., see Sablonniere, B. 183
Dempewolk, J., see Bours, J. 336
Ekundayo, O.
——, Laakso, I.,Hiltunen, R., and Seppänen, T.
Rapid analyses of chromenes (benzopyrans) by gas chromatography–mass spectrometry 358
Elenbring, K., see Hjertén, S. 47
Elhilali, M., see Bouthillier, F. 171
Ellow, L.-E., see Bergenhem, N. 383
Eng, G.
——, Johannesen, R. B., Tierney, E. J., Bellama, J. M. and Brinckman, F. E.
Holistic molecular conformation and total surface area calculations as predictors for solution properties and chromatographic parameters 1
Fanali, S.
——, Masia, P. and Ossicini, L.
Paper electrophoretic study of ion-pair formation. XVI. Resolution of optical isomers of some cobalt(III) complexes with amino acid ligands 388
Farkas, M., see Papp, O. 19
Formstecher, P., see Sablonniere, B. 183
Foster, W. S.
—— and Jarrett, H. W.
Melittin-silica, a high-pressure affinity chromatography resin for calmodulin 99
Franceschini, N., see Amicosante, G. 366
Frommer-Filep, M., see Gál, S. 217
Gál, S.
——, Tar, A., Frommer-Filep, M., Toth-Martinez, B. L., Hernádi, F. J. and Kiss, L.
Use of chromatofocusing for separation of β-lactamases. VIII. Analytical chromatofocusing of chromosomal cephalosporinases from four *Klebsiella* strains 217

García-Pérez, A.I.
——, Recio, M. N., Sancho, P. and Luque, J.
Partitioning behaviour of rat bone marrow cells in aqueous two-phase systems. Dependence of cell partition on the interfacial tension and electrical potential difference between the phases 131
Gasanova, G. Z., see Zaslavsky, B. Yu. 123
Ghosh, P.
——, Sil, P. and Thakur, S.
Spray reagent for the detection of coumarins and flavonoids on thin-layer plates 285
Giger, W., see Czuczwa, J. 233
——, see Marcomini, A. 243
Glöckner, G.
Normal- and reversed-phase separation of copolymers prepared from styrene and ethyl methacrylate 280
Godoy, A. Adaya, see Ladron de Guevara, O. 350
Green, M. R., see Haken, J. K. 145
Guieze, P., see Varotsis, N. 159
Guimbard, J. P.
——, Person, M. and Vergnaud, J. P.
Determination of residual solvents in pharmaceutical products by gas chromatography coupled to a head-space injection system and using an external standard 109
Haken, J. K.
—— and Green, M. R.
Cleavage of urea–formaldehyde resin in cross-linked alkyd systems 145
Hart, R. J., see Sworn, G. 307
Hashimoto, H., see Yamashita, J. 275
Hawthorne, S. B.
—— and Miller, D. J.
Directly coupled supercritical fluid extraction–gas chromatographic analysis of polycyclic aromatic hydrocarbons and polychorinated biphenyls from environmental solids 63
Hellberg, U., see Johansson, B.-L. 85
Hermecz, I., see Papp, O. 11, 19
Hernádi, F. J., see Gál, S. 217
Hiltunen, R., see Ekundayo, O. 358
Hjertén, S.
——, Elenbring, K., Kilár, F., Liao, J.-L., Chen, A. J. C., Siebert, C. J. and Zhu, M.-D.
Carrier-free zone electrophoresis, displacement electrophoresis and isoelectric focusing in a high-performance electrophoresis apparatus 47
Hoogmartens, J., see Cachet, Th. 343
Ishii, D., see Takeuchi, T. 324
Ishiwatari, R., see Morinaga, S. 225
Ito, Y.
Foam counter-current chromatography with the cross-axis synchronous flow-through coil planet centrifuge 77
James, L. B.
Amino acid analysis: the histidine hangover 271
Jarrett, H. W., see Foster, W. S. 99
Johannesen, R. B., see Eng, G. 1
Johansson, B.-L.
——, Hellberg, U. and Wennberg, O.
Determination of the leakage from Phenyl-Sepharose CL-4B Phenyl-Sepharose FF and Phenyl-Superose in bulk and column experiments 85
Józan, M., see Papp, O. 11
Junginger, H. E., see Roorda, W. E. 355
Kale, N. R., see Talib, Y. Y. 373
Karve, M. S.,see Talib, Y. Y. 373
Kikuchi, J.
——, Nakamura, K., Nakata, O. and Morikawa, Y.
Determination of monosaccharides constituting the glycosides in saponins by high-performance liquid chromatography 319
Kilár, F., see Hjertén, S. 47
Kiss, L., see Gál, S. 217
Kita, H., see Yamashita, J. 275
König, W. A., see Weller, O. 263
Kotzé, J. M., see Thomas, A. C. 292
Kuwamoto, T., see Okada, T. 35
Laakso, I., see Ekundayo, O. 358
Ladron de Guevara, O.
——, Adaya Godoy, A. and Cortinas de Nava, C.
Quantification of halothane, enflurane and nitrous oxide by means of a simple gas chromatographic method 350
Lawrence, J. F.
—— and Charbonneau, C. F.
Separation of sulphite adducts by ion chromatography with oxidative amperometric detection 379
Lefebvre, P., see Sablonniere, B. 183
Lehoux, J.-G., see Bouthillier, F. 171
Leuenberger, C., see Czuczwa, J. 233
Liao, J.-L., see Hjertén, S. 47
Luque, J., see García-Pérez, A. I. 131
Mahmudov, A. U., see Zaslavsky, B. Yu. 123
Marcomini, A.
——, Capri, S. and Giger, W.
Determination of linear alkylbenzenesulphonates, alkylphenol polyethoxylates and nonylphenol in waste water by high-performance liquid chromatography after enrichment on octadecylsilica 243
Marinucci, M. C., see Amicosante, G. 366
Marletta, G. Pertoldi, see Zonta, F. 207

Marrs, W. M., see Sworn, G. 307
Masia, P., see Fanali, S. 388
Matthee, F. N., see Thomas, A. C. 292
Miheeva, L. M., see Zaslavsky, B. Yu. 123
Miller, D. J., see Hawthorne, S. B. 63
Miyano, S., see Yamashita, J. 275
Morikawa, Y., see Kikuchi, J. 319
Morinaga, S.
—— and Ishiwatari, R.
Gas chromatographic determination of C_1–C_5 low-molecular-weight organic acids in alkaline permanganate oxidation products of humic substances 225
Moschidis, M. C.
—— and Andrikopoulos, N. K.
Separation of the phosphono analogues of phosphatidylcholine and phosphatidylethanolamine from related phospholipids by high-performance liquid chromatography 363
Nakamura, K., see Kikuchi, J. 319
Nakata, O., see Kikuchi, J. 319
Nava, C. Cortinas de, see Ladron de Guevara, O. 350
Numakura, T., see Yamashita, J. 275
Ōi, S., see Yamashita, J. 275
Okada, T.
Elution chracteristics of polyol–borate eluents in anion-exchange chromatography. I. Eluent strength of sugar–, uronate–borate eluents 27
—— and Kuwamoto, T.
Elution characteristics of polyol–borate eluents in anion-exchange chromatography. II. Tartrate–borate eluent for non-suppressed anion chromatography 35
Oliva, B., see Amicosante, G. 366
Oratore, A., see Amicosante, G. 366
Oshima, T., see Sagara, K. 253
Ossicini, L., see Fanali, S. 388
Papp, O.
——, Józan, M., Szász, Gy. and Hermecz, I.
Correlation between molar refraction and gas chromatographic retention data 11
——, Szász, Gy., Farkas, M., Simon, G. and Hermecz, I.
Some problems in the correlation of molecular parameters and connectivity index 19
Person, M., see Guimbard, J. P. 109
Pertoldi Marletta, G., see Zonta, F. 207
Porath, J., see Belew, M. 197
Radwan, F., see Bouthillier, F. 171
Recio, M. N., see García-Pérez, A. I. 131
Rink, H., see Bours, J. 336
Roets, E., see Cachet, Th. 343
Roorda, W. E.
——, Verbeek, J. and Junginger, H. E.
Simple gas chromatographic determination of the purity of hydroxyethyl methacrylate 355
Rosén, S., see Bergenhem, N. 383
Rubio, N.
——, Brieva, A. and Alcazar, B.
Purification of allergens by high-performance liquid chromatography. IV. Purification of the allergen of olive pollen (*Olea europea*) 312
Sablonniere, B.
——, Lefebvre, P., Formstecher, P. and Dautrevaux, M.
Improved stokes radius measurement of the glucocorticoid receptor using TSK G4000SW and TSK G3000SW high-performance size-exclusion columns. Analytical and preparative applications 183
Sagara, K.
——, Oshima, T. and Yoshida, T.
Rapid and simple determination of sennoside A and B in Rhei Rhizoma by ion-pair high-performance liquid chromatography 253
Sancho, P., see García-Pérez, A. I. 131
Schaake, S., see Bours, J. 336
Schafer, R., see Tedesco, J. L. 299
Schulze, J., see Weller, O. 263
Seppänen, T., see Ekundayo, O. 358
Siebert, C. J., see Hjertén, S. 47
Sil, P., see Ghosh, P. 285
Simon, M., see Papp, O. 19
Stancher, B., see Zonta, F. 207
Strašák, M.
—— and Bystrický, S.
Partial resolution of cobalt(III) chelate complexes by gel permeation chromatography 331
Suzuki, T., see Yamashita, J. 275
Sworn, G.
——, Marrs, W. M. and Hart, R. J.
Characterization of carrageenans by high-performance size-exclusion chromatography using a LiChrospher 1000 DIOL column 307
Szász, Gy., see Papp, O. 11, 19
Takai, N., see Yamashita, J. 275
Takeuchi, T.
—— and Ishii, D.
Micro high-performance liquid chromatography of poly(ethylene glycols) with indirect photometric detection 324
Talib, Y. Y.
——, Karve, M. S., Bhide, S. V. and Kale, N. R.
Separation of starch components based on ligand-induced adsorption of amylose on cellulose 373
Tar, A., see Gál, S. 217
Tarutani, T., see Yokoyama, T. 151

Tedesco, J. L.
—— and Schafer, R.
Increased yield of the phenylthiocarbamyl-amino acids tyrosine, cysteine and phenyl-alanine using diethylether-based liquid–liquid extraction 299
Thakur, S., see Ghosh, P. 285
Thomas, A. C.
——, Kotzé, J. M. and Matthee, F. N.
Characterization of *Botrytis* by gas–liquid chromatography 292
Tierney, E. J., see Eng, G. 1
Tizzani, M. I., see Amicosante, G. 366
Toth-Martinez, B. L., see Gál, S. 217
Tremp, J., see Czuczwa, J. 233
Vanderhaeghe, H., see Cachet, Th. 343
Varotsis, N.
—— and Guieze, P.
Microdistillation: a method combining gas and liquid chromatography to characterize petroleum liquids 159
Verbeek, J., see Roorda, W. E. 355
Vergnaud, J. P., see Guimbard, J. P. 109
Weller, O.
——, Schulze, J. and König, W. A.
Separation of racemic pharmaceuticals by high-performance liquid chromatography on silica gel modified with carbohydrate residues 263
Wennberg, O., see Johansson, B.-L. 85
Yamanaka, C., see Yokoyama, T. 151
Yamashita, J.
——, Numakura, T., Kita, H., Suzuki, T., Ōi, S., Miyano, S., Hashimoto, H. and Takai, N.
High-performance liquid chromatographic separation of enantiomers on axially chiral binaphthalene derivatives bonded to silica gel 275
Yip, T.-T., see Belew, M. 197
Yokoyama, T.
——, Yamanaka, C. and Tarutani, T.
Formation of silicato complexes of aluminium in aqueous solution 151
Yoshida, T., see Sagara, K. 253
Zaslavsky, B. Yu.
——, Miheeva, L. M., Gasanova, G. Z. and Mahmudov, A. U.
Effect of polymer composition on the relative hydrophobicity of the phases of the biphasic system aqueous dextran–poly(ethylene glycol) 123
Zhu, M.-D., see Hjertén, S. 47
Zonta, F.
——, Stancher, B. and Pertoldi Marletta, G.
Simultaneous high-performance liquid chromatographic analysis of free carotenoids and carotenoid esters 207